

# Donning and Doffing Personal Protective Equipment when undertaking **Aerosol Generating Procedure (AGPs)**

#### The following are considered AGPs

Induction of sputum

Open respiratory tract suctioning

Tracheal intubation and extubation

Awake\* bronchoscopy

Awake\* ear, nose, and throat (ENT) airway procedures that involve respiratory suctioning Awake\* upper gastro-intestinal endoscopy Cardiopulmonary resuscitation (CPR) Dental procedures (using high speed or high

frequency devices, for example ultrasonic scalers/high

speed drills)

#### The following are NOT Considered to be AGPs

Manual facemask ventilation Administration of pressurised humidified oxygen Non-invasive ventilation (NIV) including CPAP High flow nasal oxygen (HFNO)

Insertion or removal of supraglottic airways Administration of Entonox Administration of medication via nebulization Taking a nasopharyngeal swab

# **Donning PPE in isolation room with** lobby

### Don PPE in lobby:

- 1. Gown (long sleeved fluid repellent, disposable) if patient with confirmed/suspected respiratory infection and/or risk of significant bodily fluid exposure, if not, use plastic apron instead
- 2. FFP3 respirator, ensure good seal, perform personal fit check
- **3.** Eye protection (prescription glasses alone are NOT sufficient) – use visors in preference to goggles whenever available
- **4.** Gloves pull on so they go over the cuffs of the gown

### Doffing PPE in isolation room with lobby

Surgery or post-mortem procedures (like high speed

cutting / drilling) likely to produce aerosol from the

respiratory tract (upper or lower) or sinuses.

Tracheostomy procedures (insertion or removal)

(Start in patient's room standing near sink and bin, well away from patient)

- 1. Remove gloves (roll off) and discard into foot operated bin
- 2. Wash hands with soap and water, dry hands
- 3. Remove gown, roll down and try not to touch the outside, discard into foot operated bin (or remove apron)
  - 4. Remove eye protection and discard into foot operated bin
    - 5. Decontaminate hands
    - 6. Exit room into lobby
  - 7. Remove FFP3 without touching the front of the mask and discard into foot operated bin
    - 8. Wash hands with soap and water, dry hands
      - 9. Exit lobby and shut door
      - 10. Gel Hands outside the lobby

### **Donning PPE in side room without lobby** or for a cohort bay

- PPE must be put on outside the side room or
- Follow the above steps

# Doffing PPE in side room without lobby or for a cohort bay

- Follow steps 1 to 4 above
- Wash hands with soap and water, dry hands
  - Exit the side room or cohort bay
- Remove FFP3 mask and discard to designated foot operated bin
  - Decontaminate hands with gel

<sup>\*</sup>Awake includes 'conscious' sedation (excluding anaesthetised patients with secured airway).